arXiv:1202.1736v1 [hep-th] 8 Feb 2012

Comments on Correlation Functions of Large Spin Operators and Null Polygonal Wilson Loops

Carlos A. Cardona[a,b] *

[a] *Instituto de Astronomía y Física del Espacio (CONICET-UBA)*
C.C. 67 - Suc. 28, 1428 Buenos Aires, Argentina

[b] *Physics Department, University of Buenos Aires, CONICET.*
Ciudad Universitaria, 1428 Buenos Aires, Argentina

Abstract

We discuss the relation between correlation functions of twist-two large spin operators and expectation values of Wilson loops along light-like trajectories. After presenting some heuristic field theoretical arguments suggesting this relation, we compute the divergent part of the correlator in the limit of large 't Hooft coupling and large spins, using a semi-classical worldsheet which asymptotically looks like a GKP rotating string. We show this diverges as expected from the expectation value of a null Wilson loop, namely, as $(\ln\mu^{-2})^2$, μ being a cut-off of the theory.

Contents

1 Introduction **2**

2 Field Theory Considerations **4**

3 Pohlmeyer reduction for strings on AdS_3 **6**

4 Large spin limit of the GKP String in reduced fields **8**
4.1 Divergent area A_{div} 10

5 Conclusions **16**

---

* `cargicar@iafe.uba.ar`

1 Introduction

In the last ten years much progress has been made concerning single trace operators in $\mathcal{N} = 4$ SYM theory at both weak and strong coupling in the context of the AdS/CFT correspondence [1]. On one hand, the integrability of the theory in the planar limit [2]-[8] has led to substantial advance, mainly in the computation of anomalous dimensions of these operators. Indeed, it has guided us to a deeper understanding on the spectrum of conformal anomalous dimensions [9]-[12]. On the other hand, at the strong coupling limit of the theory, single-trace twist-two operators have been suggested to correspond to string states with angular momentum in the AdS_5 sector of string theory on $AdS_5 \times S^5$ [13]. Concretely, twist two operators of the form[1]

$$\mathcal{O}_S = \mathrm{Tr}(\Phi \bigtriangledown_{\{\mu_1} \ldots \bigtriangledown_{\mu_S\}} \Phi) V^{\mu_1} \ldots V^{\mu_S} \,, \tag{1.1}$$

with spin S and conformal dimension for large spin given by

$$\Delta = S + f(\lambda) \ln S \,, \tag{1.2}$$

are described at strong coupling by a macroscopic rotating string on AdS_5. At large S, the leading contribution to the anomalous dimension comes from the classical solution, whose energy satisfies the same scaling with the spin as the conformal dimension (1.2) [13, 14, 15].

Other important objects which have been extensively studied during the last years are Wilson loops made from null lines. These have several interesting properties. To mention just a few, they are related to scattering amplitudes in gauge theories [16]-[20], particularly to deep inelastic scattering amplitudes in QCD [21]. But more relevant for the purposes of this note, they can be used also to compute the anomalous dimensions of the operators we would like to consider, namely, those of the form (1.1). This has been done at weak coupling in [22, 23] and at strong coupling in [24]. The anomalous dimension 'completely' determines the two-point correlation function of the corresponding operator, suggesting that there is a relation between correlation functions of operators $\mathcal{O}_S$ and expectation values of Wilson loops, in a similar way as the relation found between the latter and correlation functions of scalars in [25]. In this note we explore some possible connections between these two objects, both at strong coupling and weak coupling, which have been suggested before in [26].

At strong coupling, the expectation value of a Wilson loop is given by the area of the worldsheet which ends on it at the boundary of AdS_5 [27]. Light-like polygonal Wilson loops have been extensively studied at strong coupling by Maldacena *et al.* [26, 28, 29, 30]. Expectation values of null Wilson loops contain ultraviolet divergences at weak and strong coupling. At weak coupling, they are associated with the fast moving particles through the Wilson light-like lines as well as with the existence of cusps, and at strong coupling, these are due to the infinite area of the world-sheet ending on the sides of the Wilson loops. On the other hand, correlators of large spin operators $\mathcal{O}_S$ have two types of divergences. One type comes from the light-like distance limit between the operator and the other one comes from the large spin limit. As we will see, at strong coupling these two are somehow related.

---

[1] In this equation $\{\cdots\}$ means totally symmetrized and V^μ is an arbitrary vector.

The calculation of the n-point function at strong coupling in the leading semiclassical approximation was shown in [31, 32, 33] to be intrinsically related to finding an appropriate classical solution of a string which ends on the boundary of AdS. Let $V_{H_i}(z_i)$, $i = 1, ..., n$ be the $n-$vertex operators inserted at points z_i on the worldsheet corresponding to a folded string.[2] For large string tension, the n-point function should be dominated by its semiclassical limit i.e. by the action evaluated at its stationary point

$$\langle \prod_{i=1}^{n} V_{H_i}(z_i) \rangle \sim e^{-\frac{R^2}{2\pi\alpha'}A} , \tag{1.3}$$

where A is the string action on $AdS_5 \times S^5$ in conformal gauge, which in the semiclassical limit is reduced to the area of the worldsheet. For the case of the two-point function, the appropriate classical solution corresponds to some analytical continuation of the GKP string [31], which reaches the boundary of AdS when the spin of the string goes to infinity. This implies that, in the limit we are considering, the area of the worldsheet diverges and should be somehow regularized, in the same way as for expectation values of Wilson lines at strong coupling. This should also happen in the classical solutions corresponding to higher point functions. Hence, the n-point functions split in a divergent contribution and a finite (regularized) part,

$$\langle \prod_{i=1}^{n} V_{H_i}(z_i) \rangle \sim e^{-\frac{R^2}{2\pi\alpha'}(A_{\rm div}+A_{\rm reg})} . \tag{1.4}$$

In this note we will study the divergent part of the correlator and we will show that it scales in the same way as the null polygonal Wilson loop associated with it[3].

The presentation is organized as follows. In section 2 we will give some hints from the field theory perspective supporting the relation between correlation functions of large spin operators (1.1) and expectation values of null polygonal Wilson loops. In section 3 we will summarize the Pohlmeyer reduction of strings on AdS_3. This is a useful technique which reduces the problem of solving the string equations of motion to that of finding a solution of a generalized Sinh-Gordon equation. This reduction is particularly powerful when we want to explore and use the integrability of string theory on AdS spaces. In section 4 we will use the Pohlmeyer reduced fields in order to study worldsheet solutions which contain asymptotically folded strings at large spin and from these we will compute the corresponding regularized area.

---

[2] We consider planar AdS/CFT duality, i.e. tree-level string theory.

[3] In a recent paper [34] the finite contribution $A_{\rm reg}$ of the 3-point correlator has been computed by exploiting the integrability of string theory on AdS_3 through the use of the Pohlmeyer reduced system.

2 Field Theory Considerations

In this section we will give some field theoretical evidence supporting the relation between the large spin limit of correlators involving the fields (1.1) and expectation values of null polygonal Wilson loops. We are considering operators of the schematic form $\mathcal{O}_S = \text{Tr}[\Phi(V^\mu \bigtriangledown_\mu)^S \Phi]$, which are characterized by the spin number S and the conformal weight Δ. In the large spin limit, we expect that each operator will emit a pair of particles and we will compute the correlation functions in the limit in which the emitted particles tend to follow light-like directions. In order to make direct contact with Wilson loops, we recall that the operator O_S arises in a power series expansion of a Wilson line

$$W(V) = \text{Tr}\left(\Phi(0)\, \mathrm{e}^{\int_0^V A_\mu dV^\mu} \Phi(V^\mu)\right) = \sum_{S=0}^{\infty} \frac{1}{S!} \mathcal{O}_S(V^\mu)\,. \tag{2.1}$$

We will choose the vector V^μ along a null direction x^- which means we are taking the Wilson line along the given null direction. That will not necessarily happen for finite spin S, but it is certainly the case for infinite spin limit. In that limit, correlators of Wilson lines of the type above should be dominated by the operators with large spin. Hence we will consider correlators of Wilson lines along null directions. Alternatively, as it has been argued in [26] and can be deduced through AdS/CFT considerations, the insertion of an operator $\mathcal{O}_S$ produces a displacement between fields Φ along the x^- direction joined by an adjoint Wilson line connecting the two-operators. Moreover, the larger the spin the larger the distance between fields and they become displaced along the null direction as the spin goes to infinity.

In this section we will follow the lines of [21] in order to compute the correlator between Wilson lines (2.1) along null directions. Let us suppose one of the line operators emits two particles of momentum p and p' as shown in figure 1[4], and compute its propagator at one loop. Consider the absorption of a gluon with momentum k and gauge potential $A^\mu(k)$ emitted by the Wilson line. In the momentum representation, this process contributes to the correlator the following vertex[5]

$$g \int \frac{d^4k}{(2\pi)^4} \frac{(2\,p' + k)_\mu}{((p'+k)^2 + i\epsilon)} A^\mu(k)\,. \tag{2.2}$$

In the massless limit or light-like trajectories, this vertex has singularities when k is collinear to p' and for soft gluons $k \sim 0$. It has been argued in [21] that when the momentum k is collinear to p', the components of $A^\mu(k)$ transverse to k contribute to higher twist and can be neglected in the limit we are considering.

---

[4] The particles p and p' are single particle states of the scalar Φ

[5] Here we have used the scalar-vector vertex, see appendix in [39] .



Fig. 1: One loop correction to the two-particles emission process from a Wilson line. Right hand side of the orange dotted line corresponds to the correction to the propagator of the particle with momentum p'.

Hence only longitudinal polarization survives and A^μ becomes pure gauge,

$$A^\mu(k) = k^\mu \left(\frac{y \cdot A(k)}{y \cdot k + i\epsilon} \right) , \tag{2.3}$$

Being y^μ a vector along the path followed by the particle. Inserting (2.3) into (2.2), the absorption vertex is now given by,

$$g \int \frac{d^4k}{(2\pi)^4} \left(\frac{y \cdot A(k)}{y \cdot k + i\epsilon} \right) = \int_0^\infty d\tau \, y_\mu A^\mu(y\tau) = \int_0^\infty dy \cdot A(y) \, . \tag{2.4}$$

Summing all contributions at any number of loops of $p'-$collinear gluons, we should finally obtain that the propagator corresponding to the emitted particle in the light-cone limit due to the interaction with gluons turns out to be

$$\mathcal{G}(\Delta x^+) = G_{\text{free}}(\Delta x^+) \mathcal{P}\text{exp} \left(ig \int_{x_1^-}^{x_2^-} dy \cdot A(y) \right) , \tag{2.5}$$

where G_{free} corresponds to the propagator for the free particle propagating between points x_1^- and x_2^- separated by a light-like distance Δx^+. This can be justified in the following way. Since the Wilson line (2.1) is dominated by the contributions of local operators at large spin S, it could be thought of as a very fast particle propagating in the x^- direction which after a given threshold starts to emit particles and gluons. When the gluons and particles emitted are collinear, a divergence in the propagator of the emitted particle occurs, dominating the whole propagator. Since collinear gluons are pure gauge, they cannot change the state of the particle but only its phase. That pure phase is given by the Wilson line (2.5) along the light-like direction x^+.

Another way to see this is based on the arguments given in [25]. The propagator of a scalar particle propagating between points x_i^- and x_{i+1}^- interacting with a gauge field is given by replacing the free scalar propagator by the propagator in the background gluon field $S(x_i^-, x_{i+1}^-; A)$, which satisfies the Green equation

$$iD^\mu D_\mu S(x, y; A) = \delta^4(x - y) \, , \tag{2.6}$$

with $D^\mu = \partial_x^\mu - ig[A^\mu(x),]$. In the light-cone limit, it is convenient to look for a solution of the above equation of the form (2.5) times some function of x which goes to one when x tends to $x^\pm$.

Concerning to divergences coming from soft glouns, we expect they cancel against virtual gluon corrections as in deep inelastic scatterings in QCD.

In principle, besides the gluon absorption vertex, we should consider scalar absorption vertices also. However it has been claimed in [23, 25], using OPE arguments, that such kind of vertices only contribute at higher twist and consequently their contributions are suppressed in the light-like limit.

Finally, putting all the Wilson lines together, i.e, replacing local operators by Wilson lines of the type (2.1) and interchanging particle propagators with (2.5), the entire correlator becomes a polygonal Wilson loop with light-like edges, which in the very large spin limit we expect to be dominated by the contributions of local operators in the expansion (2.1) with large S, in a similar way like in the expectation value of a Wilson line with a single cusp [24].

3 Pohlmeyer reduction for strings on AdS_3

In order to set some notation in this section we briefly review the Pohlmeyer reduction of strings on AdS_3 proposed in [35, 36] (see also [37, 28] and references therein). We closely follow the notation of [28]. It is worth mentioning that along this note we will consider only an AdS_3 factor of the total AdS_5 space. However, we think the results could be generalized to AdS_5.

AdS_d spaces can be written as the following hyperboloid in $R^{2,d-1}$

$$\vec{Y} \cdot \vec{Y} = -Y_{-1}^2 - Y_0^2 + Y_1^2 + ... + Y_{d-1}^2 = -1\,. \tag{3.1}$$

In terms of these embedding coordinates, the equations of motion for the bosonic string are given by

$$\partial\bar{\partial}\vec{Y} - (\partial\vec{Y} \cdot \bar{\partial}\vec{Y})\vec{Y} = 0\ , \tag{3.2}$$

and the Virasoro constraints read

$$\partial\vec{Y} \cdot \partial\vec{Y} = \bar{\partial}\vec{Y} \cdot \bar{\partial}\vec{Y} = 0\,. \tag{3.3}$$

Let us start defining the reduced fields α and p in AdS_3 as,

$$\begin{aligned} e^{2\alpha(z,\bar{z})} &= \frac{1}{2}\partial\vec{Y} \cdot \bar{\partial}\vec{Y}\ , \\ p = -\frac{1}{2}\vec{N} \cdot \partial^2\vec{Y}\ , &\qquad \bar{p} = \frac{1}{2}\vec{N} \cdot \bar{\partial}^2\vec{Y}\ , \\ N_a &= \frac{e^{-2\alpha}}{2}\epsilon_{abcd}Y^b\partial Y^c\bar{\partial}Y^d\,. \end{aligned} \tag{3.4}$$

From (3.4) and (3.2) it can be shown that $p = p(z)$ is a holomorphic function [6], where we have parametrized the world-sheet in terms of complex variables z and $\bar{z}$. Let us introduce the following basis of four-vectors in $R^{2,2}$

$$\vec{q}_1 \;=\; \vec{Y} \;, \qquad \vec{q}_2 = e^{-\alpha}\bar{\partial}\vec{Y} \;, \qquad \vec{q}_3 = e^{-\alpha}\partial\vec{Y} \;, \qquad \vec{q}_4 = \vec{N} \;, \tag{3.5}$$

which satisfies

$$\vec{q}_1^{\;2} = -1 \;, \qquad\qquad \vec{q}_2.\vec{q}_3 = 2, \qquad \vec{q}_4^{\;2} = 1 \,, \tag{3.6}$$

with the remaining $\vec{q}_i.\vec{q}_j = 0$. The last property along with the equivalence between $SO(2,2)$ and $SL(2) \times SL(2)$ allows to write the basis vectors in the following matrix representation

$$W = \frac{1}{2}\begin{pmatrix} \vec{q}_1 + \vec{q}_4 & \vec{q}_2 \\ \vec{q}_3 & \vec{q}_1 - \vec{q}_4 \end{pmatrix} , \tag{3.7}$$

W being an $SL(2) \times SL(2)$ element. The components of this matrix have indices $W_{\alpha\dot{\alpha},a\dot{a}}$. The first two indices denote rows and columns in the above matrix, while the other two are associated with the space-time bispinor representation of each $\vec{q}_i$, i.e,

$$(q_i)_{a\dot{a}} = q_{i,\mu}\sigma^{\mu}_{a\dot{a}}, \qquad \sigma^{\mu} = (1, i\sigma_3, \sigma_1, -\sigma_2)\,. \tag{3.8}$$

The two $SL(2)$ symmetries of the AdS_3 target space corresponds to the $SL(2)$ group acting on the index a and the $SL(2)$ that acts on the index $\dot{a}$. The basis vectors (3.5) define the $SL(2)$ connections $B^{L,R}$ given by [36]

$$B^L_z = \begin{pmatrix} \frac{1}{2}\partial\alpha & -e^{\alpha} \\ -e^{-\alpha}p(z) & -\frac{1}{2}\partial\alpha \end{pmatrix}, \qquad B^L_{\bar{z}} = \begin{pmatrix} -\frac{1}{2}\bar{\partial}\alpha & -e^{-\alpha}\bar{p}(\bar{z}) \\ -e^{\alpha} & \frac{1}{2}\bar{\partial}\alpha \end{pmatrix} \tag{3.9}$$

$$B^R_z = \begin{pmatrix} -\frac{1}{2}\partial\alpha & e^{-\alpha}p(z) \\ -e^{\alpha} & \frac{1}{2}\partial\alpha \end{pmatrix}, \qquad B^R_{\bar{z}} = \begin{pmatrix} \frac{1}{2}\bar{\partial}\alpha & -e^{\alpha} \\ e^{-\alpha}\bar{p}(\bar{z}) & -\frac{1}{2}\bar{\partial}\alpha \end{pmatrix} . \tag{3.10}$$

The consistency conditions of the equations (3.2) imply that these connections are flat

$$\partial B^L_{\bar{z}} - \bar{\partial}B^L_z + [B^L_z, B^L_{\bar{z}}] = 0 \;, \quad \partial B^R_{\bar{z}} - \bar{\partial}B^R_z + [B^R_z, B^R_{\bar{z}}] = 0\,. \tag{3.11}$$

These imply that p is a holomorphic function and that α satisfies the generalized sinh-Gordon equation

$$\partial\bar{\partial}\alpha(z,\bar{z}) - e^{2\alpha(z,\bar{z})} + |p(z)|^2 e^{-2\alpha(z,\bar{z})} = 0\,. \tag{3.12}$$

We are especially interested in the area of the worldsheet, which is given in terms of the reduced fields by,

$$A = 4\int d^2z e^{2\alpha}\,. \tag{3.13}$$

Given a solution of the generalized sinh-Gordon model, one can find the associated classical string worldsheet. In order to do that, consider the auxiliary linear problem [28]

$$\partial\psi^L_{\alpha} + (B^L_z)_{\alpha}^{\;\beta}\psi^L_{\beta} = 0, \qquad \bar{\partial}\psi^L_{\alpha} + (B^L_{\bar{z}})_{\alpha}^{\;\beta}\psi^L_{\beta} = 0\,,$$

---

[6] For the real solutions considered in this paper, $p(z)$ and $\bar{p}(\bar{z})$ are complex conjugates. This condition could in principle be relaxed.

$$\partial\psi^R_{\dot\alpha} + (B^R_z)_{\dot\alpha}{}^{\dot\beta}\psi^R_{\dot\beta} = 0, \qquad \bar\partial\psi^R_{\dot\alpha} + (B^R_z)_{\dot\alpha}{}^{\dot\beta}\psi^R_{\dot\beta} = 0\,. \tag{3.14}$$

We denote by $\psi^L_{\alpha,a}$, $a = 1,2$, and $\psi^R_{\dot\alpha,\dot a}$, $\dot a = 1,2$, the two independent solutions for the left and right linear equations respectively. Since the connections are in $SL(2)$, we can define an $SL(2)$ invariant product and use it to normalize the pair of solutions as

$$\langle\psi^L_a,\,\psi^L_b\rangle \equiv \epsilon^{\beta\alpha}\psi^L_{\alpha,a}\psi^L_{\beta,b} = \epsilon_{ab}, \qquad \langle\psi^R_{\dot a},\,\psi^R_{\dot b}\rangle \equiv \epsilon^{\dot\beta\dot\alpha}\psi^R_{\dot\alpha,\dot a}\psi^R_{\dot\beta,\dot b} = \epsilon_{\dot a\dot b}\,. \tag{3.15}$$

It is easy to see from (3.14) that the normalizations (3.15) are both constant and then they can be evaluated at any point. Now we can reconstruct the space-time worlsheet coordinates from these solutions through the matrix $W_{\alpha\dot\alpha,a\dot a}$, as has been explained in [28]

$$W_{\alpha\dot\alpha,a\dot a} = \psi^L_{\alpha,a}\psi^R_{\dot\alpha,\dot a}\,. \tag{3.16}$$

This could be justified noticing that each component of W in (3.7) is a null vector in $R^{2,2}$, so they can be written as a product of spinors. The explicit form of the target coordinates is given by the element q_1 in (3.7), which can be written as

$$Y_{a\dot a} = \begin{pmatrix} Y_{-1}+Y_2 & Y_1-Y_0 \\ Y_1+Y_0 & Y_{-1}-Y_2 \end{pmatrix}_{a,\dot a} = \psi^L_{\alpha,a}\psi^R_{\dot\beta,\dot a}\ , \tag{3.17}$$

and similarly for the other q_i.

4 Large spin limit of the GKP String in reduced fields

In this section we would like to explain how to compute the divergent part of the worldsheet area $A_{\rm div}$ following an approach similar to the one used in [28]. In order to do that we will use the reduced fields just described in the section above in the particular case of a worldsheet which contains asymptotic large spin limit of GKP strings.

In global coordinates the GKP string at large spin S is given in worldsheet coordinates $(\tau,\,\sigma)$ by [38, 13]

$$t = \kappa\tau, \quad \theta = \omega\tau, \quad \rho = \rho(\sigma), \quad \omega \to \kappa \sim \frac{1}{\pi}\ln S\,. \tag{4.1}$$

In embedding coordinates its looks like

$$Y_{-1} + iY_0 = \mathrm{e}^{\kappa\tau}\cosh\rho(\sigma), \quad Y_1 + iY_2 = \mathrm{e}^{-\kappa\tau}\sinh\rho(\sigma)\,. \tag{4.2}$$

Inserting the last coordinates in (3.4), we get that the GKP strings are described by the following reduced fields in cylindrical coordinates $w = \tau + i\sigma$,

$$p(w) = -\frac{\kappa^2}{4}, \quad \mathrm{e}^{2\alpha(w,\bar w)} = \frac{\kappa^2}{4}\,. \tag{4.3}$$

Going to the plane through the conformal transformation $z = \mathrm{e}^w$ we have

$$p(z) = -\frac{\kappa^2}{4\,z^2}, \quad \mathrm{e}^{2\alpha(w,\bar w)} = \sqrt{p\bar p}\,. \tag{4.4}$$

As we can see, $p(z)$ encodes the information on the vertex positions, which in this case appear at $z = 0, \infty$. Now we would like to solve the linear problem (3.14) associated to the folded string. It is easier to solve it first in w-coordinates, since the reduced fields do not depend on the coordinates there. Doing so we obtain (see [34])

$$\tilde{\psi} = \mathcal{A}\psi = \frac{1}{\sqrt{2}}\, \mathrm{e}^{\pm\frac{i\kappa}{2}(\zeta^{-1}w - \zeta\bar{w})} \begin{pmatrix} 1 \\ \pm 1 \end{pmatrix} , \tag{4.5}$$

where $\mathcal{A} = \mathrm{diag}(p^{-1/4}\,\mathrm{e}^{\alpha/2}, p^{1/4}\,\mathrm{e}^{-\alpha/2})$ and we have introduced a spectral parameter ζ in order to write the solutions in a compact way. The actual space time solutions are given by $\zeta = 1$ for ψ^L and $\zeta = i$ for ψ^R. Taking $z = r\,\mathrm{e}^{i\phi}$, on the $z-$plane, the above solutions behave as

$$\eta^{L\pm} \sim \mathrm{e}^{\pm\kappa\phi}\, v_{L\pm}, \quad \eta^{R\pm} \sim r^{\pm\kappa}\, v_{R\pm} \,, \tag{4.6}$$

where $v's$ are given by

$$v_{L+} = \begin{pmatrix} -i \\ 1 \end{pmatrix}, \quad v_{L-} = \begin{pmatrix} -1 \\ i \end{pmatrix}, \quad v_{R+} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad v_{R-} = \begin{pmatrix} -1 \\ 1 \end{pmatrix} . \tag{4.7}$$

Every solution can be written as a linear combination of the above η solutions. Hence, we choose the following arbitrary combinations as a basis of solutions

$$\psi^L_a = c_{l,a}\eta^{L,l}, \quad \psi^R_{\dot{a}} = c_{l,\dot{a}}\eta^{R,l}, \quad a, \dot{a} = 1, 2 \,. \tag{4.8}$$

Generically, we are looking for solutions behaving as those above near each singularity produced by the insertion of a vertex operator. Namely, generically we assume the function $p(z)$ has n-singularities at points z_i, describing the insertion of vertex operators in the $z-$plane. More explicitly, we assume the function $p(z)$ behaves near singularities in the following way

$$p(z) \sim -\sum_{i=1}^{n} \frac{\kappa_i^2}{4(z - z_i)^2} \,, \tag{4.9}$$

κ_i being a parameter related to the large spin of the operator in the corresponding singularity, α is given by $\mathrm{e}^{2\alpha(w,\bar{w})} = \sqrt{p\bar{p}}$ near each insertion, and we assume the singularities are far enough from each other in order to preserve the approximation. Hence, near each singular point, the basis of solutions behaves as

$$\psi^L_{a\,i} \sim c_{l,a\,i}\,\eta^{L,l}, \quad \psi^R_{\dot{a}\,i} \sim c_{l,\dot{a}\,i}\,\eta^{R,l}, \quad a, \dot{a} = 1, 2, \quad i = 1, ..., n-1 \,. \tag{4.10}$$

In order to see how the solutions approach each vertex insertion in the generic case, it is convenient to write them in $(z, \bar{z})$ coordinates

$$\eta^{R\,\pm} \sim \prod_{i=1}^{n} [(z - z_i)(\bar{z} - \bar{z}_i)]^{\pm k_i/2}, \quad \eta^{L\,\pm} \sim \prod_{i=1}^{n} \left(\frac{z - z_i}{\bar{z} - \bar{z}_i}\right)^{\pm k_i/2} . \tag{4.11}$$

The target space coordinates can be recovered from the above solutions using (3.17), such that in Poincare coordinates (Z, x_μ) we have

$$\frac{1}{Z} = Y_{1\dot{1}}, \quad x^+ = \frac{Y_{1\dot{2}}}{Y_{1\dot{1}}}, \quad x^- = \frac{Y_{2\dot{1}}}{Y_{1\dot{1}}}, \quad x^\pm = x_0 \pm x_1 \,. \tag{4.12}$$

As we see from (4.11), some solutions get bigger and some get smaller when we approach the vertex. Therefore the target coordinates will be dominated by the big solutions when evaluated near the insertions. Then, from (3.17) and (4.10), they are schematically approximated near the singularities by

$$Y_{a\dot{a}} = c^{L\,\mathrm{big}}_{a\,i} c^{R\,\mathrm{big}}_{\dot{a}\,i} f_{\mathrm{big}}(z - z_i, \bar{z} - \bar{z}_i)\,, \tag{4.13}$$

where the label 'big' on the coefficients means we keep only those multiplying the $\eta^{R,L}$ which become bigger near the given z_i and $f_{\mathrm{big}}(z - z_i, \bar{z} - \bar{z}_i)$ is the large contribution coming from the $\eta^{R,L}$ which grows up the most.

$$\frac{1}{Z} = c^{L\,\mathrm{big}}_{1\,i} c^{R\,\mathrm{big}}_{\dot{1}\,i} f_{\mathrm{big}}(z - z_i, \bar{z} - \bar{z}_i)\,, \quad x^{+} = \frac{c^{R\,\mathrm{big}}_{\dot{2}\,i}}{c^{R\,\mathrm{big}}_{\dot{1}\,i}}\,, \quad x^{-} = \frac{c^{L\,\mathrm{big}}_{2\,i}}{c^{L\,\mathrm{big}}_{1\,i}}\,. \tag{4.14}$$

4.1 Divergent area A_{div}

As we already mentioned, the source of divergences for the area comes from the regions close to the insertions, which are precisely the regions where the string approaches the boundary of AdS. Therefore, it is obvious that putting a worldsheet cutoff near each singularity is completely equivalent to putting a cutoff in target space for the coordinate Z close to the boundary, which at the same time corresponds, through AdS/CFT, to putting an ultraviolet cutoff on the energy of the process. In this section we will show how the divergent contribution to the correlator (1.4) at strong coupling, A_{div}, is related to the space-time cutoff and how it depends on the dynamics of the string at the boundary.

From equations (4.6) and (4.10) we can see the large spin solutions diverge near $\phi = \pm\pi/2$, $r = \pm\infty$, and therefore, the worldsheet is approaching the boundary at those points. Moreover, as we will see, the worldsheet approaches the boundary on light-like trajectories. This can be seen from equation (4.14), realizing that some coefficients $c^{\mathrm{big}}_{a,i}$ are equal for consecutive i, i.e, the points i and $i+1$ have the same, lets say, x^{+}_{i} coordinate and therefore they are joined by a null line.

The area is given by (3.13), and the divergent part of it comes from the regions near singularities, and as we said, near each singularity $\mathrm{e}^{2\alpha} \sim \sqrt{p\bar{p}}$, hence

$$A_{\mathrm{div}} = 4 \int d^2\, z\, \sqrt{p(z)\bar{p}(\bar{z})}\,. \tag{4.15}$$

But since the leading contribution to the divergent area comes from the regions very near the singularities of $p(z)$, we are going to isolate the contributions of each z_i and approximate A_{div} as

$$A_{\mathrm{div}} = -4 \sum_{i=1}^{n} \int_{|z-z_i|>\epsilon_i} d^2\, z\, \frac{k_i^2}{|z - z_i|^2} \sim -4\pi \sum_{i=1}^{n} k_i^2 \ln\epsilon_i\,, \tag{4.16}$$

where we have regularized the integral through a worldsheet cutoff ϵ. As we mentioned, when we approach the singular points z_i, the worldsheet gets closer to the boundary, and hence the worldsheet cutoff ϵ should be related to some physical cutoff $1/Z = 1/\mu$, $\mu << 1$, which corresponds to putting a brane very close to the boundary where the tips of the folded string will end in the limit of infinite spin.

Without loss of generality, let us start considering only three operator insertions, taking κ_1, $\kappa_3 > 0$ and $\kappa_2 < 0$. As we see from equation (4.13), the behaviour of the target coordinates near the boundary is well approximated by the big solutions near each vertex. Which solution becomes big depends on the radial distance to the insertion as well as on the angle around the insertion. In order to visualize the behavior of the target space coordinates near the insertions, we display the figures 2 and 3. Here we will consider only the half $z-$plane because in the large S limit, the folded strings bend into two equal pieces, and then the area could be computed as two times the area corresponding to a half of the folded worldsheet (for details see [34]).



fig.2: Half $z-$plane with three holes of size ϵ representing the position of the insertions.

Points 1, 3, 5 correspond to angles $\pi/2$ and points 2, 4, 6 correspond to angles $-\pi/2$ around the corresponding insertion. As we will see, red lines and green lines map to the null directions defining the Wilson loop. Therefore, the points $1, 2, ..., 6$ correspond in the space-time to the cusp of the null Wilson loop. Let us denote the positions of those cusps as $(Z_l,\, x_l^{\pm})$, and see how we approach the boundary around each cusp. Since κ_1, $\kappa_3 > 0$ and $\kappa_2 < 0$, we have according to (4.13),

$$\begin{aligned}
\frac{1}{Z_1} &= c^{L+}_{1\,1} c^{R-}_{\dot{1}\,1} \left(\frac{z-z_1}{\bar{z}-\bar{z}_1}\right)^{k_1/2} |z-z_1|^{-k_1}, \quad x_1^+ = \frac{c^{R-}_{\dot{2}\,1}}{c^{R-}_{\dot{1}\,1}}, \quad x_1^- = \frac{c^{L+}_{2\,1}}{c^{L+}_{1\,1}}, \\
\frac{1}{Z_2} &= c^{L-}_{1\,1} c^{R-}_{\dot{1}\,1} \left(\frac{z-z_1}{\bar{z}-\bar{z}_1}\right)^{-k_1/2} |z-z_1|^{-k_1}, \quad x_2^+ = \frac{c^{R-}_{\dot{2}\,1}}{c^{R-}_{\dot{1}\,1}}, \quad x_2^- = \frac{c^{L-}_{2\,1}}{c^{L-}_{1\,1}}, \\
\frac{1}{Z_3} &= c^{L+}_{1\,2} c^{R+}_{\dot{1}\,2} \left(\frac{z-z_2}{\bar{z}-\bar{z}_2}\right)^{k_2/2} |z-z_2|^{-k_2}, \quad x_3^+ = \frac{c^{R+}_{\dot{2}\,2}}{c^{R+}_{\dot{1}\,2}}, \quad x_3^- = \frac{c^{L+}_{2\,2}}{c^{L+}_{1\,2}}, \\
\frac{1}{Z_4} &= c^{L-}_{1\,2} c^{R+}_{\dot{1}\,2} \left(\frac{z-z_2}{\bar{z}-\bar{z}_2}\right)^{-k_2/2} |z-z_2|^{-k_2}, \quad x_4^+ = \frac{c^{R+}_{\dot{2}\,2}}{c^{R+}_{\dot{1}\,2}}, \quad x_4^- = \frac{c^{L-}_{2\,2}}{c^{L-}_{1\,2}}, \\
\frac{1}{Z_5} &= c^{L+}_{1\,3} c^{R-}_{\dot{1}\,3} \left(\frac{z-z_3}{\bar{z}-\bar{z}_3}\right)^{k_3/2} |z-z_3|^{-k_3}, \quad x_5^+ = \frac{c^{R-}_{\dot{2}\,3}}{c^{R-}_{\dot{1}\,3}}, \quad x_5^- = \frac{c^{L+}_{2\,3}}{c^{L+}_{1\,3}}, \\
\frac{1}{Z_6} &= c^{L-}_{1\,3} c^{R-}_{\dot{1}\,3} \left(\frac{z-z_3}{\bar{z}-\bar{z}_3}\right)^{-k_3/2} |z-z_3|^{-k_3}, \quad x_6^+ = \frac{c^{R-}_{\dot{2}\,3}}{c^{R-}_{\dot{1}\,3}}, \quad x_6^- = \frac{c^{L-}_{2\,3}}{c^{L-}_{1\,3}}.
\end{aligned} \tag{4.17}$$

As we mentioned in section 4, we can see from the equations above that the target positions of points connecting red lines on the worldsheet (see fig.2) live on the same null lines in space-time, i.e $x_1^+ = x_2^+$, $x_3^+ = x_4^+$, $x_5^+ = x_6^+$.



Fig. 2: Mapping of the boundaries of the half-worlsheet into space-time light-like lines.

The coordinates displayed in (4.17) are good approximations to the exact string solution only in the neighborhood of each insertion (think about each insertion as an asymptotic state in target space). Each asymptotic state is connected to each other through the worldsheet, but since we do not know the exact solution, we cannot determine how these states are exactly correlated. One way to relate each insertion to each other is by using the so-called monodromy matrix. As the auxiliary linear problem has two independent solutions, with a basis given by (ψ_1, ψ_2), we expect that as one analytically continues them around a singularity, they get linearly transformed as

$$\begin{pmatrix} \psi_1' \\ \psi_2' \end{pmatrix} = S \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} . \tag{4.18}$$

It is worth mentioning that the monodromy matrix should belong to $SL(2,R)$,[7] and therefore it satisfies $\det|S| = 1$. Applying (4.18) to (4.10), we see that we can reinterpret (4.18) as

$$c_{a,i}^{L\alpha} = S_{\beta}^{L\alpha} c_{a,i}^{L\beta} , \tag{4.19}$$

with a similar relation for the c^R coefficients and α, $\beta = \pm$.

---

[7] We have written $\vec{Y} \in$SO(2,2) as an element $Y_{a\dot{a}} \sim \psi_a^L \psi_{\dot{a}}^R$ of SL(2,R)×SL(2,R), or in other words, the eigenvectors of S should live on a representation of SL(2,R).

According to the discussion in section 2, we expect that the leading contribution to the correlator comes from the particles propagating between the operators through nearly null trajectories. Let us start imposing a null trajectory condition to the path connecting cusp two to cusp three

$$x_2^- = \frac{c_{2,1}^{L-}}{c_{1,1}^{L-}} \equiv x_3^- = \frac{c_{2,2}^{L+}}{c_{1,2}^{L+}} = \frac{S_{\beta,\,2\to1}^{L+} c_{2,1}^{L\beta}}{S_{\beta,\,2\to1}^{L+} c_{1,1}^{L-}}\,, \tag{4.20}$$

therefore, the S^L-matrix should satisfy $S_{+,\,2\to1}^{L+} = 0$, and from the determinant we get $S_{-,\,2\to1}^{L+} S_{+,\,2\to1}^{L-} = -1$, i.e

$$S_{2\to1}^L = \begin{pmatrix} 0 & \pm1 \\ \mp1 & \gamma_{2\to1}^L \end{pmatrix}\,, \tag{4.21}$$

where we have defined $S_{-,\,2\to1}^{L-} = \gamma_{2\to1}^L$. Doing the same for the points connecting cusps 4 and 5 and cusps 6 and 1, we end up with[8]

$$S_{3\to2}^L = \begin{pmatrix} 0 & \pm1 \\ \mp1 & \gamma_{3\to2}^L \end{pmatrix}\,, \quad S_{1\to3}^L = \begin{pmatrix} 0 & \pm1 \\ \mp1 & \gamma_{1\to3}^L \end{pmatrix}\,. \tag{4.22}$$

Now we would like to compute the distances between consecutive points, or the separation length of points living on a null line. Let us start considering

$$x_1^- - x_2^- = \frac{c_{2,1}^{L+} c_{1,1}^{L-} - c_{1,1}^{L+} c_{2,1}^{L-}}{c_{1,1}^{L+} c_{1,1}^{L-}} = -\frac{\langle \psi_{1,1}^L\,,\,\psi_{2,1}^L \rangle}{c_{1,1}^{L+} c_{1,1}^{L-}}\,. \tag{4.23}$$

Recalling we have normalized the basis of solutions in each sector as $\langle \psi_{1,i}^L\,,\,\psi_{2,i}^L \rangle = 1$, we get

$$x_1^- - x_2^- = -\frac{1}{c_{1,1}^{L+} c_{1,1}^{L-}} \equiv x_{12}^-\,. \tag{4.24}$$

On the other hand, we have $x_2^- \equiv x_3^-$, and then

$$x_2^+ - x_3^+ = \frac{c_{\dot2,1}^{R-}}{c_{\dot1,1}^{R-}} - \frac{c_{\dot2,2}^{R+}}{c_{\dot1,2}^{R+}} \equiv \frac{a_{\dot2,\dot1}^R}{c_{\dot1,1}^{R-} c_{\dot1,2}^{R+}}\,, \tag{4.25}$$

where we have defined

$$\begin{aligned} a_{\dot2,\dot1}^R &= c_{\dot2,1}^{R-}(S_{+\,2\to1}^{R+} c_{\dot1,1}^{R+} + S_{-\,2\to1}^{R+} c_{\dot1,1}^{R-}) - c_{\dot1,1}^{R-}(S_{+\,2\to1}^{R+} c_{\dot2,1}^{R+} + S_{-\,2\to1}^{R+} c_{\dot2,1}^{R-}) \\ &= \langle \psi_{1,1}^R\,,\,\psi_{2,1}^R \rangle\, S_{+\,2\to1}^{R+} = \gamma_{2\to1}^R\,, \end{aligned} \tag{4.26}$$

with $S_{+\,2\to1}^{R+} = \gamma_{2\to1}^R$. The parameters $\gamma_{i+1\to i}^{R,L}$ parametrize our ignorance on the exact solution and we cannot compute them only using the approximated solutions around the insertions. Introducing a cutoff μ_i given by the value of Z_i closest to the boundary and recalling that the worldsheet cutoff is defined by $|z - z_i| > \epsilon_i$ we have from (4.17)

---

[8] These monodromy matrices take the form of the Stokes matrices considered in [28] by using a different argument.

$$- \ln\mu_1 = \ln(c^{L+}_{1,1} c^{R-}_{\dot{1},1}) - \kappa_1 \ln\epsilon_1 \,, \tag{4.27}$$

$$- \ln\mu_2 = \ln(c^{L-}_{1,1} c^{R-}_{\dot{1},1}) - \kappa_1 \ln\epsilon_2 \,. \tag{4.28}$$

Where we have dropped the term depending on ϕ because the cutoff has been defined for the entire angular region around the insertion. Summing up (4.27) and (4.28) and using (4.24), we get

$$- \kappa_1 \ln\epsilon_1\epsilon_2 = (- \ln\mu_1\mu_2 + \ln x^-_{12} + 2 \ln c^{R,-}_{\dot{1},1}) \,. \tag{4.29}$$

In the strict infinite spin limit, the parameter κ goes to infinity as well, as can be seen from (4.1), introducing another kind of divergence, different from that associated to the worldsheet approaching the boundary and therefore independent of the worldsheet cutoff. In order to isolate the dependence of κ on the space-time cutoff, we evaluate the first equation in (4.17) a slightly away from the insertions but not too far in order to avoid spoiling the approximation, and find

$$g(z^h_1)\kappa_1 = - \ln\mu_1 - \ln(c^{L+}_{1,1} c^{R-}_{\dot{1},1}) = - \ln\mu_1 + \ln u_1 \,, \tag{4.30}$$

where the point z^h_1 is arbitrary, not too far from the insertion and we have defined $u_1 = c^{L+}_{1,1} c^{R-}_{\dot{1},1}$. The function $g(z)$ is given by,

$$g(z = r\, \mathrm{e}^{i\phi}) = \ln \frac{\mathrm{e}^{\phi}}{r} \,. \tag{4.31}$$

Let us argue about the natural choice of z^h_1. For $r_i < 1$ the space-time coordinates start to feel the presence of the divergence associated with the proximity of the boundary and for $r_i > 1$ we start to spoil the approximation. On the other hand for ϕ close to $\pm\pi/2$ the coordinates are close to the boundary again. Therefore, the natural choice seems to be $z_i = \mathrm{e}^{i\pi}$ and then $g(z^h_i) = \pi$.[9]
Inserting (4.30) into (4.29) we have that

$$- \pi\kappa_1^2 \ln\epsilon_1\epsilon_2 = (- \ln\mu_1 + \ln u_1)(- \ln\mu_1\mu_2 + \ln x^-_{12} + 2 \ln c^{R,-}_{\dot{1},1}) \,. \tag{4.32}$$

Similarly, we get

$$\begin{aligned} - \pi\kappa_2^2 \ln\epsilon_2\epsilon_3 &= (- \ln\mu_2 + \ln u_2)(- \ln\mu_2\mu_3 + \ln x^-_{34} + 2 \ln c^{R,+}_{\dot{1},2}) \,, && (4.33) \\ -\pi\kappa_3^2 \ln\epsilon_3\epsilon_1 &= (- \ln\mu_3 + \ln u_3)(- \ln\mu_3\mu_1 + \ln x^-_{56} + 2 \ln c^{R,-}_{\dot{1},3}) \,. && (4.34) \end{aligned}$$

Of course we should take the same cutoff in different directions, i.e. $\mu_i = \mu$ and $\epsilon_i = \epsilon$, but the quantities u_i, $i = 1, 2, 3$ do not have to be equal in different directions. However, we will take them approximately equal in comparison with the $\ln\mu$ accompanying them $u_1 \sim u_2 \sim u_3$, hence

---

[9] This argument seems to be a bit arbitrary, but we mainly would like to show the logarithmic scaling of κ with μ.

$$\begin{aligned} A_{\rm div} &\sim -4\pi \sum_{i=1}^{3} k_i^2 \ln\epsilon_i \\ &\sim 4(-\ln\mu + \ln u_1)\Big(\sum_{i=1}^{3} 2\ln\tilde{c}_{1,i}^{R,-} - 6\ln\mu + \ln x_{12}^- + \ln x_{34}^- + \ln x_{56}^-\Big)\,, \end{aligned} \tag{4.35}$$

where we have defined $\tilde{c}_{1,1}^{R,-} = c_{1,1}^{R,-}$, $\tilde{c}_{1,2}^{R,-} = c_{1,2}^{R,+}$, $\tilde{c}_{1,3}^{R,-} = c_{1,3}^{R,-}$. Using (4.17) and (4.26) plus the similar relations for x_{45}^+ and x_{61}^+, it is easy to see that

$$\sum_{i=1}^{3} 2\ln\tilde{c}_{1,i}^{R,-} = \ln x_{23}^+ + \ln x_{45}^+ + \ln x_{61}^+ + \ln\left(\gamma_{2\to1}^R \gamma_{3\to2}^R \gamma_{1\to3}^R\right)\,. \tag{4.36}$$

Putting all those things together we get

$$\begin{aligned} A_{\rm div} \sim\ & 4(-\ln\mu + \ln u_1)\times \\ & \big(-6\ln\mu + \ln x_{12}^- + \ln x_{34}^- + \ln x_{56}^- + \ln x_{23}^+ \\ & + \ln x_{45}^+ + \ln x_{61}^+ + \ln\left(\gamma_{2\to1}^R \gamma_{3\to2}^R \gamma_{1\to3}^R\right)\big)\,. \end{aligned} \tag{4.37}$$

For the sake of simplicity, let us redefine variables in the following way

$$x_{12}^- \equiv \mathbf{x}_1^-\,,\ \ x_{34}^- \equiv \mathbf{x}_2^-\,,\ \ x_{56}^- \equiv \mathbf{x}_3^-\,,\ \ x_{23}^+ \equiv \mathbf{x}_1^+,\ \ x_{45}^+ \equiv \mathbf{x}_2^+,\ \ x_{61}^+ \equiv \mathbf{x}_3^+\,. \tag{4.38}$$

We can now rewrite the divergent area as [10]

$$A_{\rm div} \sim 4\sum_{i=1}^{3}(-\ln\mu + \ln u_i)(-2\ln\mu + \ln[\mathbf{x}_i^+\mathbf{x}_i^-] + \ln\left(\gamma_{2\to1}^R \gamma_{3\to2}^R \gamma_{1\to3}^R\right))\,. \tag{4.39}$$

As we have seen, the coefficients $c_{a,i}^{L\alpha}$ are associated with the space-time coordinates. So we would like to write the expression only in terms of the x^+, x^-. We illustrate that by working on the first term from the sum (4.39),

$$A_{\rm div} \sim 4\sum_{i=1}^{3}(-\ln\mu + \ln u_i)B_i\,, \tag{4.40}$$

being,

$$B_i \equiv (-2\ln\mu + \ln[\mathbf{x}_i^+\mathbf{x}_i^-] + \ln\left(\gamma_{2\to1}^R \gamma_{3\to2}^R \gamma_{1\to3}^R\right))\,. \tag{4.41}$$

Summing and subtracting $(-\ln\mu + \ln c_{1,1}^{L-} c_{1,2}^{R+})$ inside the first bracket in $A_{\rm div}$ we get,

$$(-2\ln\mu + \ln x_{12}^- + \ln x_{23}^+)B_1 - (-\ln\mu + \ln\tilde{u}_1)B_1\,, \tag{4.42}$$

---

[10] Notice that with (4.38) we are associating two consecutive light-like directions for each i, ($i = 1, 2, 3$), hence in the expression for $A_{\rm div}$ (4.39) or (4.43) the number of light-like segments involved are 2×3 and then (4.43) should be related to the hexagonal Wilson loop.

where we have defined $\tilde{u}_i = c^{L-}_{1,1} c^{R+}_{1,2}$. Doing the same for the remaining terms into the sum (i.e, summing and subtracting the lacking factor that builds the corresponding coordinates), we end up with,

$$A_{\text{div}} \sim 4\sum_{i=1}^{3}\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]\right)^2$$
$$+4\sum_{i=1}^{3}\left[\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]\right)\ln\left(\gamma^R_{2\to1}\gamma^R_{3\to2}\gamma^R_{1\to3}\right)+\left(\ln\frac{\mu}{\tilde{u}_i}\right)\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]+\ln\left(\gamma^R_{2\to1}\gamma^R_{3\to2}\gamma^R_{1\to3}\right)\right)\right]. \tag{4.43}$$

Defining $\tilde{\gamma}^R_{i\to i+1} \equiv \gamma^R_{i+1\to i}$, we generalize A_{div} to the case of $2n-$null lines or $n-$vertex insertions as,

$$A_{\text{div}} \sim 4\sum_{i=1}^{n}\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]\right)^2$$
$$+4\sum_{i=1}^{n}\left[\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]\right)\ln\left(\prod_{i=1}^{n+1}\tilde{\gamma}^R_{i\to i+1}\right)+\left(\ln\frac{\mu}{\tilde{u}_i}\right)\left(\ln[\frac{\mathbf{x}_i^+\mathbf{x}_i^-}{\mu^2}]+\ln\left(\prod_{i=1}^{n+1}\tilde{\gamma}^R_{i\to i+1}\right)\right)\right]. \tag{4.44}$$

Where $n+1=1$. The form of the first line coincides with the most divergent term in the expectation value of a null Wilson loop, computed at strong coupling in [28], and given the presence of the matrices $\tilde{\gamma}^R$, the two terms in the second line will depend on the exact solution.

5 Conclusions

In this paper we have considered a light-like limit of correlation functions involving operators of high spin. We have argued that in the large spin limit, correlators should be related to expectation values of Wilson loops along paths lying on the light cone. At strong 't Hooft coupling we performed a semi-classical computation of the divergent part of the correlator and found that it scales with the cut-off in the same way the expectation value of a null Wilson loop does.

In order to confirm the correspondence between Wilson loops and high spin correlation functions, at least at semi-classical level, we should find an exact classical solution of a string in AdS which contains GKP classical states and has the topology of a sphere with n-vertex insertions. As far as we know, that solution is unknown. However, the correlator could be computed semi-classically by using integrability, as has been done at strong coupling for the finite part in [34] and at weak coupling in [40]. The next interesting step could be to use the results in [34] in order to extract the finite part of the Wilson loops from the correlators.

Acknowledgments

I am grateful to Martin Kruczenski and Juan Maldacena for very useful discussions and interesting comments and suggestions. I also thank Carmen Nuñez who encouraged me to publish this note and for carefully read the manuscript. I would also like to thank Oscar Bedoya and Humberto Gomez for reading the first version of this note. I also like to thank to the Institute for Advanced Studies at Princeton for hospitality during the early stages of this work.

This work was supported by CONICET PIP 112-200801-00507, MINCYT (Ministerio de Ciencia, Tecnología Innovación Productiva de Argentina), University of Buenos Aires UBACyT X161.

References

[1] J. M. Maldacena, "The Large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. **2** (1998) 231 [Int. J. Theor. Phys. **38** (1999) 1113] [hep-th/9711200].

[2] I. Bena, J. Polchinski and R. Roiban, "Hidden symmetries of the $AdS_5 \times S^5$ superstring," Phys. Rev. **D 69** (2004) 046002 [hep-th/0305116].

[3] J.A. Minahan and K. Zarembo, " The Bethe-ansatz for N = 4 super Yang-Mills," JHEP **03** (2003) 013 [hep-th/0212208].

[4] V.A. Kazakov, A. Marshakov, J.A. Minahan and K. Zarembo, " Classical / quantum integrability in AdS/CFT," JHEP **05** (2004) 024 [hep-th/0402207].

[5] V. A. Kazakov and K. Zarembo, "Classical / quantum integrability in non-compact sector of AdS/CFT," JHEP **0410**, 060 (2004) [arXiv:hep-th/0410105].

[6] N. Beisert, V. Dippel and M. Staudacher, "A novel long range spin chain and planar N = 4 super Yang-Mills," JHEP **07** (2004) 075 [hep-th/0405001].

[7] G. Arutyunov, S. Frolov and M. Staudacher, "Bethe ansatz for quantum strings," JHEP *10* (2004) 016 [hep-th/0406256].

[8] N. Beisert and M. Staudacher, " Long-range PSU(2,2—4) Bethe ansaetze for gauge theory and strings," Nucl. Phys. **B 727** (2005) 1 [hep-th/0504190].

[9] N. Gromov, V. Kazakov and P. Vieira, " Exact Spectrum of Anomalous Dimensions of Planar $N = 4$ Supersymmetric Yang-Mills Theory," Phys. Rev. Lett. **103** (2009) 131601 [arXiv:0901.3753].

[10] D. Bombardelli, D. Fioravanti and R. Tateo, " Thermodynamic Bethe Ansatz for planar AdS/CFT: a proposal," J. Phys. A 42 (2009) 375401 [arXiv:0902.3930].

[11] N. Gromov, V. Kazakov, A. Kozak and P. Vieira, “Exact Spectrum of Anomalous Dimensions of Planar N = 4 Supersymmetric Yang-Mills Theory: TBA and excited states,” Lett. Math. Phys. **91** (2010) 265 [arXiv:0902.4458].

[12] G. Arutyunov and S. Frolov, “Thermodynamic Bethe Ansatz for the $AdS_5 \times S^5$ Mirror Model,” JHEP **05** (2009) 068 [arXiv:0903.0141].

[13] S. S. Gubser, I. R. Klebanov and A. M. Polyakov “A semi-classical limit of the Gauge/string correspondence,” [arXiv:hep-th/0204051].

[14] M. Kruczenski, “Spiky strings and single trace operators in gauge theories,” JHEP **0508** (2005) 014 [hep-th/0410226].

[15] A. Jevicki, K. Jin, C. Kalousios and A. Volovich,
“Generating AdS String Solutions,” JHEP **0803** (2008) 032 [arXiv:0712.1193 [hep-th]].

[16] L. F. Alday and J. M. Maldacena, “Gluon scattering amplitudes at strong coupling,” JHEP **0706**, 064 (2007) [arXiv:0705.0303 [hep-th]].

[17] L. F. Alday and R. Roiban, “Scattering Amplitudes, Wilson Loops and the String/Gauge Theory Correspondence,” Phys. Rept. **468**, 153 (2008) [arXiv:0807.1889 [hep-th]].

[18] J. M. Drummond, G. P. Korchemsky and E. Sokatchev, “Conformal properties of four-gluon planar amplitudes and Wilson loops,” Nucl. Phys. B **795**, 385 (2008) [arXiv:0707.0243 [hep-th]].

[19] A. Brandhuber, P. Heslop and G. Travaglini, “MHV Amplitudes in N=4 Super Yang-Mills and Wilson Loops,” Nucl. Phys. B **794**, 231 (2008) [arXiv:0707.1153 [hep-th]].

[20] J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “Conformal Ward identities for Wilson loops and a test of the duality with gluon amplitudes,” arXiv:0712.1223 [hep-th].

[21] G. P. Korchemsky and G. Marchesini, “Structure function for large x and renormalization of Wilson loop,” Nucl. Phys. B **406**, 225 (1993) [hep-ph/9210281].

[22] G.P. Korchemsky, “Asymptotics of the Altarelli-Parisi-Lipatov evolution kernels of parton distributions,” Mod. Phys. Lett. A**4** ,1257 (1989)

[23] G.P. Korchemsky, G. Marchesini, “Structure functions for large x and normalization of Wilson loops,” Nucl. Phys. B **406**, 225 (1993) [hep-ph/9210281].

[24] M. Kruczenski, “A Note on twist two operators in N=4 SYM and Wilson loops in Minkowski signature,” JHEP **0212**, 024 (2002) [hep-th/0210115].

[25] L. F. Alday, B. Eden, G. P. Korchemsky, J. Maldacena and E. Sokatchev, “From correlation functions to Wilson loops,” JHEP **1109**, 123 (2011) [arXiv:1007.3243 [hep-th]].

[26] L. F. Alday, D. Gaiotto, J. Maldacena, A. Sever and P. Vieira, “An Operator Product Expansion for Polygonal null Wilson Loops,” JHEP **1104**, 088 (2011) [arXiv:1006.2788 [hep-th]].

[27] J. M. Maldacena, “Wilson loops in large N field theories,” Phys. Rev. Lett. **80**, 4859 (1998) [hep-th/9803002].

[28] L. F. Alday and J. Maldacena, “Null polygonal Wilson loops and minimal surfaces in Anti-de-Sitter space,” JHEP **0911**, 082 (2009) [arXiv:0904.0663 [hep-th]].

[29] L. F. Alday, D. Gaiotto and J. Maldacena, “Thermodynamic Bubble Ansatz,” arXiv:0911.4708 [hep-th].

[30] L. F. Alday, J. Maldacena, A. Sever and P. Vieira, “Y-system for Scattering Amplitudes,” J. Phys. A A **43**, 485401 (2010) [arXiv:1002.2459 [hep-th]].

[31] A. A. Tseytlin, “On semiclassical approximation and spinning string vertex operators in AdS”, Nucl. Phys. B **664**, 247 (2003) [hep-th/0304139].

[32] E. I. Buchbinder, “Energy-Spin Trajectories in $AdS_5 \times S^5$ from Semiclassical Vertex Operators,” JHEP **1004**, 107 (2010). [arXiv:1002.1716].

[33] E. I. Buchbinder and A. A. Tseytlin, “On semiclassical approximation for correlators of closed string vertex operators in AdS/CFT,” JHEP **1008**, 057 (2010). [arXiv:1005.4516].

[34] Y. Kazama, S. Komatsu, “On holographic three point functions for GKP strings from integrability,” [arXiv:1110.3949 [hep-th]].

[35] K. Pohlmeyer, “Integrable Hamiltonian Systems And Interactions Through Quadratic Constraints,” Commun. Math. Phys. **46**, 207 (1976).

[36] H. J. De Vega and N. G. Sanchez, “Exact Integrability Of Strings In D-Dimensional De Sitter Space-Time,” Phys. Rev. D **47**, 3394 (1993).

[37] A. Jevicki, K. Jin, C. Kalousios and A. Volovich, JHEP **0803**, 032 (2008). [arXiv:0712.1193 [hep-th]].

[38] R. Roiban and A. A. Tseytlin, “On semiclassical computation of 3-point functions of closed string vertex operators in $AdS_5 x S^5$,” Phys. Rev. D **82**, 106011 (2010) [arXiv:1008.4921 [hep-th]].

[39] Ta-Pei Cheng and Ling-Fong Li, “Gauge theory of elementary particle physics,” Oxford University Press, 1984.

[40] J. Escobedo, N. Gromov, A. Sever and P. Vieira, “Tailoring Three-Point Functions and Integrability,” JHEP **1109**, 028 (2011) [arXiv:1012.2475 [hep-th]].